

HAND & PLASTIC SURGERY CENTRE, P.L.C.

Ronald D. Ford, M.D., F.A.C.S.
William T. Cullen, M.D., F.A.C.S.
Paul A. Luce, M.D.
Shannon D. Armstrong, M.D.
Matthew D. Martin, M.D.

Surgery of the Hand & Upper Extremity
Microvascular Surgery
Plastic & Reconstructive Surgery

245 Cherry Street, S.E., Suite 302
Grand Rapids, MI 49503
Phone (616) 459-4131 Fax (616) 459-6030

2373 64th Street, Suite 2700
Byron Center, MI 49315
Phone (616) 301-0240 Fax (616) 301-0250

THIGH LIFT POST-OP INSTRUCTIONS

AT HOME AFTER THIGH LIFT SURGERY:

- It is normal to experience bruising, swelling, numbness, soreness, tightness, burning, and drainage from your incisions.
- Swelling may occur for approximately 1-6 months, and numbness will continue to improve as healing progresses.
- You may help lessen bruising, swelling and discomfort with the use of cold compresses. For the first 72 hours, sleep with pillows under your legs to keep them in an elevated position.
- Full leg extension may be impossible for the first week. You may begin walking when you are comfortable. Don't overdo it, use your discomfort as a guide.
- Please ask your physician before you begin taking aspirin or other anti-inflammatory medications, as these can create greater risk for bleeding.

OTHER POST-OPERATIVE INSTRUCTIONS:

- Your dressings will consist of a compression garment to help reduce swelling and bruising. The sutures will be dissolvable so nothing will need to be removed.
- You may need to wait 2-3 days before you shower. At this time you may shower if you are comfortable, but leave the paper tapes over your incisions in place. Wash the area of sutures gently but thoroughly each time. You may need to apply gauze over your incisions to absorb any drainage and replace as needed. Tub bathing is not recommended until all incisions have healed.
- Your support garments will need to be worn for 4-6 weeks following surgery. You may adjust the garment as needed. It should be snug, but not uncomfortable.
- Please be careful when shaving, as your incisions may be puckered and while you are still experiencing numbness.
- Avoid sun exposure to new incision areas. These areas are very sensitive to sunlight and can be easily burned for up to a year following surgery. We recommend sunscreen to all healed incisions.
- Light exercises such as walking can begin when you are comfortable, but avoid any strenuous activities such as lifting, straining and stretching your legs.
- Strenuous activities may be gradually increased 2-3 weeks following surgery, and you may typically resume normal activities in about 4-6 weeks.
- Returning to work will vary depending on the job type. Generally you will be able to return to work approximately 7-10 days following your surgery. You may discuss this issue with your physician at the time of your post-op visit.
- Scarring will mature with time; it may take up to 3 months to a year for scars to fade and flatten.

Please contact the office immediately if any of the following occur **(616) 459-4131**:

- Fever of 100.5 or greater
- Unusual swelling, redness, bleeding or increase pain
- Persistent vomiting
- Develop hives, diarrhea, or other reactions to medicine
- Any other questions or concerns

POST-OP APPOINTMENT: _____

PRESCRIBED MEDICATION(S): _____

Please take Narcotic/OTC medication as directed and as needed. **DO NOT** drive or operate machinery if you are taking a narcotic. If you are prescribed antibiotics, take as directed until gone. Taking an incomplete course can lead to recurrence of infection. Please take antibiotics with food as this may cause upset stomach. Please ask physician before you begin taking aspirin, motrin/ibuprofen or other anti-inflammatory medications, as these can increase risk of bleeding.

****IF NAUSEATED, TRY NON-ACIDIC LIQUIDS, DRY TOAST OR OTHER BLAND FOODS****

Patient Signature: _____ **Date:** _____

Hospital Staff Signature: _____ **Date:** _____

Physician Signature: _____ **Date:** _____